


This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently amended) ~~A salt~~ Salts of a saturated, partially or fully unsaturated, heterocyclic ~~eations~~ cation ~~having the and~~ bis(trifluoromethyl)imide anion, $N(CF_3)_2^-$, which have the general formula (1)



where

X = N, P, O or S

n = an integer selected from 0, 1 or 2 ~~in such a way that~~ wherein X is saturated in accordance with its valency increased by 1,

A = a saturated, partially or fully unsaturated 3- to 8-membered hydrocarbon chain,
in which all carbon atoms apart from one may be replaced by identical or different heteroatoms selected from N, P, O and S,
where the carbon atoms of the hydrocarbon chain and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,
straight-chain or branched alkyl having 1-20 carbon atoms,
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds,
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds,
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon

atoms,
 halogen, ~~in particular~~ fluorine or chlorine, with the proviso that, for X = N,
 O, S, there is no halogen-heteroatom bond,
 -NO₂, with the proviso that there is no bond to a positively charged
 heteroatom,
 -CN, with the proviso that there is no bond to a positively charged
 heteroatom,


where the R² and/or R¹ in different and/or identical position of the
 heterocyclic ring are in each case identical or different,

where the R² and/or R¹ may be connected to one another in pairs by a
 single or double bond,

where one or more R² and/or R¹ may be partially or fully substituted by
 halogens, ~~in particular F and/or Cl~~, or partially by -CN or -NO₂, with the
 proviso that not all R² and R¹ are fully halogenated,


and where one or two carbon atoms of the R¹ and/or R² may be replaced by
 heteroatoms and/or atomic groups selected from the group -O-, -C(O)-,
 C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and
 -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅,
 where the α -position of the R¹ is not replaced for X = O, S.

2. (Currently amended) A salt ~~Salts~~ according to Claim 1, in which A is a 4-, 5- or
 6-membered hydrocarbon chain.
3. (Currently amended) A salt ~~Salts~~ according to Claim 1 in which A is a hydrocarbon
 chain in which zero, one or two carbon atoms are replaced by heteroatoms selected
 from N, P, O and S.
4. (Currently amended) A salt ~~Salts~~ according to Claim 1, in which
 R¹, R², independently of one another, have the meaning
 -H, with the proviso that there is no bond to the positively charged heteroatom,
 halogen, in particular fluorine, with the proviso that for X = N, O, S, there is no
 halogen-heteroatom bond,
 straight-chain or branched alkyl having 1-6 carbon atoms, ~~in particular CH₃, C₂H₅, n-~~



straight-chain or branched, partially or perfluorinated alkyl having 1-6 carbon atoms, in particular CF_3 , C_2F_5 , C_4F_9 .

5. (Currently amended) ~~A salt~~ Salts according to Claim 1, in which the saturated, partially or fully unsaturated, heterocyclic cation is selected from the following group:



where

X = N, P, O or S


Y = in each case, independently of one another, N, P, O, S or C,
where at least one Y = C

n = 0 for unsaturated X = O, S
1 for saturated X = O, S or for unsaturated X = N, P
2 for saturated X = N, P

m = 0 for saturated Y = O, S or for unsaturated Y = N, P
1 for saturated Y = N, P or for Y = $\text{sp}^2\text{-C}$
2 for Y = $\text{sp}^3\text{-C}$

where the radicals R^1 , R^2 are as defined in Claim 1.

6. (Currently amended) ~~A salt~~ Salts according to Claim 1, in which the saturated, partially or fully unsaturated, heterocyclic cation is selected from the following group:



where

X = in each case, independently of one another, N, P, O, S or C,
where at least one X = N, P, O or S


n = 0 for saturated X = O, S or for unsaturated X = N, P
1 for saturated X = N, P or for X = sp²-C
or for substitution on the sp² ring carbon atom
2 for X = sp³-C
or for substitution on the sp³ ring carbon atom

where n is increased by 1 for one X = N, P, O or S

A' = saturated, partially or fully unsaturated 2- to 7-membered hydrocarbon chain,
where the carbon atoms of the hydrocarbon chain are saturated by substituents R² in accordance with their valency

where the radicals R¹, R² are as defined in Claim 1, with the proviso that the substituents R¹ in the α-position to the positively charged heteroatom do not have a methylene group directly adjacent to the heterocyclic ring.

7. (Currently amended) A salt ~~Salts~~ according to Claim 1, in which the saturated, partially or fully unsaturated, heterocyclic cation has the following structure



where

X = N or P

Y = N, P, O or S

n = 0 for Y = O, S
1 for Y = N, P


A' = saturated, partially or fully unsaturated 2- to 7-membered hydrocarbon chain,

B' = saturated, partially or fully unsaturated 1- to 6-membered hydrocarbon chain,

where all carbon atoms of the hydrocarbon chains A' and B' apart from one may be replaced by identical or different heteroatoms selected from N, P, O and S and where the carbon atoms of the hydrocarbon chains A' and B' are saturated by substituents R² in accordance with their valency

and where the radicals R¹, R² are as defined in Claim 1.

8. (Currently amended) A salt ~~Salts~~ of saturated, partially or fully unsaturated, heterocyclic di-cations having the bis(trifluoromethyl)imide anion, N(CF₃)₂⁻, which have the general formula (2)



where

X, Y = each, independently of one another, N, P, O or S

n, m = an integer selected from 0, 1 or 2 in such a way that X and Y are each saturated in accordance with their valency increased by 1,

A'', B'' = saturated, partially or fully unsaturated 0- to 4-membered hydrocarbon chain,
in which the carbon atoms may be replaced by identical or different heteroatoms selected from N, P, O and S,
where at least one carbon atom is present in the chains A'' and B'' together and
where the carbon atoms of the hydrocarbon chains A'' and B'' and the heteroatoms present therein are saturated by substituents R² in accordance with their valency,

R¹, R², R³ = -H, with the proviso that there is no bond to the positively charged heteroatom,
straight-chain or branched alkyl having 1-20 carbon atoms,
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds,
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds,
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms,
halogen, in particular fluorine or chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
-NO₂, with the proviso that there is no bond to a positively charged heteroatom,
-CN, with the proviso that there is no bond to a positively charged heteroatom,

where the R¹, R² and/or R³ in different and/or identical position of the

heterocyclic ring are in each case identical or different,

where the R^1 , R^2 and/or R^3 may be connected to one another in pairs by a single or double bond,


where one or more R^1 , R^2 and/or R^3 may be partially or fully substituted by a halogen, ~~halogens, in particular~~ -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that that not all R^1 , R^2 and R^3 are fully halogenated,

and where one or two carbon atoms of the R^1 , R^2 and/or R^3 may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α -positions of the R^1 and of the R^3 are not replaced for X = O, S or Y = O, S,

where the heterocyclic di-cation is a 4-, 5-, 6-, 7-, 8- or 9-membered ring.

9. (Currently amended) A salt ~~Salts~~ according to Claim 8, in which the heterocyclic di-cation is a 5-, 6- or 7-membered ring.

10. (Currently amended) A salt ~~Salts~~ of saturated, partially or fully unsaturated, heterocyclic cations having the bis(trifluoromethyl)imide anion, $N(CF_3)_2^-$, which have the ~~general~~ formula (3)



where

X, Y = each, independently of one another, N, P, O or S

$n =$ an integer selected from 0, 1 or 2 in such a way that X is saturated in accordance with its valency increased by 1,

$m =$ an integer selected from 0, 1 or 2 in such a way that Y is saturated in accordance with its valency,

$R^1, R^2 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,
straight-chain or branched alkyl having 1-20 carbon atoms
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms,
halogen, ~~in particular~~ chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
-OR, with the proviso that the substituted heteroatom is not O or S,

where the R^2 and/or R^1 in different and/or identical position of the heterocyclic ring are in each case identical or different,


where the R^2 and/or R^1 may be connected to one another in pairs by a single or double bond,

where one or more R^2 and/or R^1 may be partially or fully substituted by a halogen, ~~halogens, in particular~~ -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that not all R^2 and R^1 are fully halogenated,

and where one or two carbon atoms of the R^1 and/or R^2 may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α -position of the R^1 is not replaced for X = O, S.


11. (Currently Amended) ~~Process~~ A process for the preparation of salts according to Claim 1, ~~characterised in that~~ wherein an alkali metal fluoride of the general formula

DF, where D selected from the group of the alkali metals, is reacted in a polar organic solvent with



where $R^F = F$ or $C_p F_{2p+1}$, where $p = 1 - 8$,

and a salt of the general formula (4)



where

$E^- = F^-, Cl^-, Br^-, I^-, BF_4^-, ClO_4^-, AsF_6^-, SbF_6^-, SbCl_6^-, PF_6^-, R^F SO_3^-, FSO_3^-, (R^F)_2 P(O)O^-, R^F P(O)_2 O^-, RSO_3^-, ROSO_3^-, \frac{1}{2} SO_3^{2-}, CN^-, SCN^-, R^F C(O)O^-, RC(O)O^-, 2,4\text{-dinitrophenolate or } 2,4,6\text{-trinitrophenolate}$, where R^F is a perfluorinated C_1 to C_8 -alkyl group or perfluorinated aryl group and R is a C_1 to C_8 -alkyl group or aryl group

$X = N, P, O \text{ or } S$

$n =$ an integer selected from 0, 1 or 2 in such a way that X is saturated in accordance with its valency increased by 1,

$A =$ a saturated, partially or fully unsaturated 3- to 8-membered hydrocarbon chain,
in which all carbon atoms apart from one may be replaced by identical or different heteroatoms selected from N, P, O and S,
where the carbon atoms of the hydrocarbon chain and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2 = -H$, with the proviso that there is no bond to the positively charged

heteroatom,
 straight-chain or branched alkyl having 1-20 carbon atoms
 straight-chain or branched alkenyl having 2-20 carbon atoms and one or
 more double bonds
 straight-chain or branched alkynyl having 2-20 carbon atoms and one or
 more triple bonds
 saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
 halogen, in particular fluorine or chlorine, with the proviso that, for X = N,
 O, S, there is no halogen-heteroatom bond,
 -NO₂, with the proviso that there is no bond to a positively charged
 heteroatom,
 -CN, with the proviso that there is no bond to a positively charged
 heteroatom,


where the R² and/or R¹ in different and/or identical position of the
 heterocyclic ring are in each case identical or different,

where the R² and/or R¹ may be connected to one another in pairs by a
 single or double bond,

where one or more R² and/or R¹ may be partially or fully substituted by a
 halogen halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂,
 with the proviso that not all R² and R¹ are fully halogenated,


and where one or two carbon atoms of the R¹ and/or R² may be replaced by
 heteroatoms and/or atomic groups selected from the group -O-, -C(O)-,
 C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and
 -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅,
 where the α-position of the R¹ is not replaced for X = O, S.

12. (Currently Amended) ~~Process~~ A process for the preparation of salts according to
 Claim 8, ~~characterised in that wherein~~ an alkali metal fluoride of the general formula
 DF, where D selected from the group of the alkali metals, is reacted in a polar organic
 solvent with



where $R^F = F$ or C_pF_{2p+1} , where $p = 1 - 8$,

and a salt of the general formula (5)



where

$E^- =$ F^- , Cl^- , Br^- , I^- , BF_4^- , ClO_4^- , AsF_6^- , SbF_6^- , $SbCl_6^-$, PF_6^- , $R^F SO_3^-$, FSO_3^- , $(R^F)_2 P(O)O^-$, $R^F P(O)_2 O^-$, RSO_3^- , $ROSO_3^-$, $\frac{1}{2} SO_3^{2-}$, CN^- , SCN^- , $R^F C(O)O^-$, $RC(O)O^-$, 2,4-dinitrophenolate or 2,4,6-trinitrophenolate, where R^F is a perfluorinated C_1 to C_8 -alkyl group or perfluorinated aryl group and R is a C_1 to C_8 -alkyl group or aryl group

$X, Y =$ each, independently of one another, N, P, O or S

$n, m =$ an integer selected from 0, 1 or 2 in such a way that X, Y are each saturated in accordance with their valency increased by 1,

$A'', B'' =$ saturated, partially or fully unsaturated 0- to 4-membered hydrocarbon chain,
in which the carbon atoms may be replaced by identical or different heteroatoms selected from N, P, O and S,
where at least one carbon atom is present in the chains A'' and B'' together and
where the carbon atoms of the hydrocarbon chains A'' and B'' and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2, R^3 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,

straight-chain or branched alkyl having 1-20 carbon atoms
 straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
 straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds
 saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
 halogen, in particular fluorine or chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
 halogen, in particular fluorine or chlorine, with the proviso that there is no halogen-heteroatom bond,
 -NO₂, with the proviso that there is no bond to a positively charged heteroatom,
 -CN, with the proviso that there is no bond to a positively charged heteroatom,

where the R¹, R² and/or R³ in different and/or identical position of the heterocyclic ring are in each case identical or different,

where the R¹, R² and/or R³ may be connected to one another in pairs by a single or double bond,

where one or more R¹, R² and/or R³ may be partially or fully substituted by halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that not all R¹, R² and R³ are fully halogenated,

and where one or two carbon atoms of the R¹, R² and/or R³ may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α-position of the R¹ is not replaced for X = O, S,

where the heterocyclic di-cation is a 4-, 5-, 6-, 7-, 8- or 9-membered ring.

13. (Currently Amended) ~~Process A~~ process according to Claim 11, ~~characterised in that wherein~~ the alkali metal fluoride employed is KF or RbF.

14. (Currently Amended) ~~Process~~ A process according to Claim 11, ~~characterised in that wherein~~ the reaction takes place at temperatures between -40°C and 80°C, in particular at 0°C to 40°C.
15. (Currently Amended) ~~Process~~ A process according to Claim 11, ~~characterised in that wherein~~ the reaction takes place in a polar organic solvent selected from the group acetonitrile, dimethoxyethane, dimethylformamide and propionitrile.
16. (Currently Amended) ~~Process~~ A process according to Claim 11, characterised in that the reaction is carried out as a one-pot reaction.
17. (Currently Amended) ~~Process~~ A process according to Claim 11, ~~characterised in that wherein~~ the reaction for $E^- = F^-$ is carried out without added alkali metal fluoride DF.
18. (Currently Amended) ~~Process~~ A process according to Claim 11, ~~characterised in that wherein~~ the starting materials for the reaction are employed in approximately equimolar ratio to one another.
19. (Currently Amended) ~~Use of a salt according to Claim 1 as~~ An ionic liquid comprising a salt according to Claim 1.
20. (Currently Amended) ~~Use of a salt according to Claim 1 as~~ An non-aqueous electrolyte comprising a salt according to Claim 1..
21. (Currently Amended) ~~Use of a salt according to Claim 1 as~~ A reagent for the introduction of $N(CF_3)_2$ groups comprising a salt according to Claim 1.
22. (Currently Amended) ~~Use of a salt according to Claim 1 as~~ A phase-transfer catalyst comprising a salt according to Claim 1.
23. (Currently Amended) ~~Use of a salt according to Claim 1 as~~ An intermediate for the synthesis of liquid-crystal compounds or active ingredients, in particular for medicaments or crop protection agents comprising a salt according to Claim 1.
24. (New) A medicaments or crop-protection agent comprising a salt according to claim 1.

25. (New) A salt according to claim 1, where one or more R^2 and/or R^1 may be partially or fully substituted by -F and/or -Cl.

26. (New) A salt according to claim 4, in which R^1 or R^2 are $-CH_3$, $-C_2H_5$, $-n-C_3H_7$, $-CH(CH_3)_2$, $-n-C_4H_9$, $-n-C_6H_{13}$, $-CF_3$, $-C_2F_5$, or $-C_4F_9$.